

# Respiratory Function in Normal Singleton Pregnancy, Multifetal Gestation and Polyhydramnios

<sup>1</sup>Shyamal Dasgupta, <sup>2</sup>Gayetri Koppolu, <sup>3</sup>Tarun Kumar Ghosh, <sup>4</sup>Susmita Kundu  
<sup>5</sup>Amitava Mukherjee, <sup>6</sup>Subhankar Dasgupta

<sup>1</sup>Assistant Professor, Department of Obstetrics and Gynecology, Institute of Postgraduate Medical Education and Research Kolkata, West Bengal, India

<sup>2</sup>Postgraduate Trainee, Department of Obstetrics and Gynecology, Institute of Postgraduate Medical Education and Research Kolkata, West Bengal, India

<sup>3</sup>Professor and Head, Department of Obstetrics and Gynecology, Institute of Postgraduate Medical Education and Research Kolkata, West Bengal, India

<sup>4</sup>Associate Professor, Department of Respiratory Medicine, Institute of Postgraduate Medical Education and Research Kolkata, West Bengal, India

<sup>5</sup>Associate Professor, Department of Obstetrics and Gynecology, Institute of Postgraduate Medical Education and Research Kolkata, West Bengal, India

<sup>6</sup>Senior Resident, Department of Obstetrics and Gynecology, Institute of Postgraduate Medical Education and Research Kolkata, West Bengal, India

**Correspondence:** Shyamal Dasgupta, Assistant Professor, Department of Obstetrics and Gynecology, 31/25-A, Mall Road Kolkata-700080, West Bengal, India, Phone: (033)2559-5702, 9433493752, e-mail: shyamaldasgupta2008@yahoo.com

## ABSTRACT

Study of respiratory functions of four groups of women: Nonpregnant, early pregnancy, singleton pregnancy at 36 weeks and twin/polyhydramnios at 36 weeks reveal that inspiratory capacity (IC), tidal volume (TV) and expiratory reserve volume (ERV) in different group pairing have statistically significant difference. This is very important during the assessment of pregnant mother suffering from respiratory or cardiopulmonary disorder.

*Objectives:* This study aims to observe the additional demands placed on the maternal respiratory system in cases of advanced normal pregnancy or over distended uterus, like multifetal gestation and polyhydramnios, and whether the average Indian woman is able to cope with these respiratory changes.

*Methods:* Around 80 women were divided into four equal groups which included non-pregnant controls, normal early pregnancy (< 20 weeks), normal singleton pregnancy at 36 weeks and multifetal gestation/polyhydramnios, underwent spirometry for static lung volume measurement. The overall comparison was done by using the Kruskal-Wallis test whereas the individual comparison by Mann-Whitney test. Final statistical computing was performed with SPSS statistical package.

*Results:* The groups were comparable on the basis of age, height and hemoglobin levels. Comparisons of various groups show statistically significant difference in tidal volume, inspiratory capacity and expiratory reserve volume specially in advance pregnancy with control nonpregnant mothers without significant adverse effect on respiratory system.

*Conclusions:* Results of the study indicate that though there are significant respiratory functional changes, healthy women, even having twin/polyhydramnios, can tolerate this well. But this knowledge is very important to assess and to formulate management of pregnant women with respiratory or cardiopulmonary compromise.

**Keywords:** Singleton pregnancy, Twin polyhydramnios, Tidal volume, Inspiratory capacity, Expiratory reserve volume.

## INTRODUCTION

The study of anatomic and physiologic adaptation of respiratory system in pregnancy is necessary for proper diagnosis and appropriate intervention in case of respiratory pathology associated with pregnancy.<sup>1</sup> Estrogen causes hyperemia, hypersecretion and mucosal edema leading to nasal obstruction specially in third trimester.<sup>2</sup> On the other hand, progesterone resets chemoreceptors at a lower partial pressure of carbon dioxide leading to slight increase in partial pressure of oxygen in arterial blood (PaO<sub>2</sub>) and decrease that of carbon dioxide (PaCO<sub>2</sub>). It results in a state of respiratory alkalosis which is compensated for increased excretion of bicarbonate via urine to maintain normal acid base balance.<sup>3</sup> Respiration in pregnancy

is more diaphragmatic where diaphragm is elevated by 4 cm. With compensatory increase in the transverse and antero-posterior diameters of chest caused by hormonal effects that loosen ligaments.<sup>4</sup> It has been found that tidal volume (TV) increases progressively throughout pregnancy by increased excursion of diaphragm<sup>5,6</sup> but inspiratory capacity (IC) and vital capacity (VC) remain almost unchanged in pregnancy.<sup>7</sup> Decreased expiratory reserve volume (ERV) and functional residual capacity (FRC) cause lungs to empty near completely during expiration leading to low defense against development of hypoxia.<sup>3</sup> Respiratory rate is unchanged, so the increased oxygen demand is met by increased depth of respiration and increased diaphragmatic excursion. Assisted reproduction

increases the incidence of multifetal gestations which impose greater oxygen demand. On the other hand, these cases along with conditions like polyhydramnios causes more diaphragmatic elevation with much relaxed ligamentous attachment leading to more pronounced respiratory changes. This study aims to observe the additional demands of oxygen on above-mentioned cases, and whether the average Indian mother is able to cope with these respiratory changes.

## METHODS

A cross-sectional study was done from 1st June 2008 to 31st May 2009, at Institute of Postgraduate Medical Education and Research (IPGMER), Kolkata, India, in the Department of Obstetrics and Gynecology with the help of Department of Respiratory Medicine. All women recruited were in the age group of 18 to 31 years, of moderate income group, mostly housewives with sedentary life style or limited physical activity. All women are healthy having no current or past history of any cardiorespiratory illness, nonsmoker and not taking any medication other than iron and calcium that is believed to alter respiratory function. Women with acute pregnancy complications, like pre-eclampsia, acute polyhydramnios, were excluded from this study. Four groups were studied: Non-pregnant, normal singleton pregnancy < 20 weeks, Normal singleton pregnancy at 36 weeks, Multifetal pregnancy and/or polyhydramnios at 36 weeks, each comprising 20 subjects. Detailed history, physical examination and baseline investigations were done for each subject to rule out any cardiorespiratory illness and anemia. All women underwent spirometry at resting state, preferably few hours after meal. Tidal volume (TV), inspiratory capacity (IC), inspiratory reserve volume (IRV), expiratory reserve volume (ERV) and vital capacity (VC) were measured on a system incorporated precision heated pneumatochograph manufactured by Medi Soft, Belgium, with proper calibration according to the ATS guidelines, and a racial correction factor of 0.85 is incorporated for Indian subjects. Parameters of lung functions were measured thrice and the best values were taken.

Statistical methods adopted in the study. Since the sample size in this study is small and hence may not follow a normal distribution, comparison may be made using nonparametric methods. The overall comparison was done by using the Kruskal-Wallis test. For individual comparison, Mann-Whitney test was applied. Statistical computing was performed with SPSS statistical package. Assuming that tidal volume is the primary variable of interest, it was calculated that a minimum of 17 subjects in each groups were required in order to detect a difference of 100 ml in TV with 80% power and 5% probability of type one error.

## RESULTS

The four groups were comparable on the basis of age, height, weight, blood pressure and hemoglobin levels. Except weight other parameters had no significant differences (Table 1).

Weight gain during pregnancy is not significant for lung function testing as standard ATS guidelines for lung function test in March 1991 was based on height, age, sex and race of the individual.

In the present study, TV progressively increased with advancement of pregnancy (mean 0.56 to 0.88 lt,  $p = 0.002$ ) but IC showed an increase specially in latter weeks (mean 1.65 to 1.95 lt,  $p = 0.016$ ). On the other hand, ERV started falling in the later weeks of pregnancy (mean from 0.76 to 0.49 lt,  $p = 0.000$ ). As the gravid uterus enlarges, pushing the diaphragm up, VC and IRV showed insignificant difference during the course of pregnancy (Table 2). Same table shows no significant difference of lung function test between normal singleton pregnancy and multifetal pregnancy/polyhydramnios group.

The comparison between various groups pairing showed that the change of TV of groups 0-2, 0-3 and 1-3 show significant statistical importance ( $p$ -values 0.015, 0.000 and 0.044 respectively). The statistical analysis of ERV of group pairing 0-2, 0-3, 1-2 and 1-3 also shows statistically important decrease ( $p$ -values 0.000, 0.000, 0.011 and 0.011 respectively). The IC in different groups pairing of 0-1, 0-3 and 1-3 also reveals statistically important increase ( $p$ -values 0.023, 0.007 and 0.034) (Table 3).

## DISCUSSION

In our study, TV increased steadily during pregnancy due to the effect of progesterone causing hyperventilation (increase depth of breathing) but the respiratory rate is constant which is corroborative to other study.<sup>3</sup> Hence, the tidal volume and subsequently minute ventilation increased as reported by other authors.<sup>2,4</sup>

The ERV is reduced by 35.8% in this study mostly in the later part of pregnancy ( $p$ -value 0.000). As the anatomical dead space and, hence, residual volume do not alter in pregnancy, the ERV is reduced due to elevated diaphragm and enlarging uterus consistent to other studies.<sup>5</sup> Inspiratory capacity is increased specially in the later part of pregnancy ( $p$ -value 0.016) mainly due to the increased tidal volume since IRV remains unchanged. These results are consistent with many other studies on respiratory changes in abnormal singleton pregnancy as mentioned above. But, presence of cardiological illness (e.g. mitral valve disease) or respiratory compromise (e.g. bronchial asthma), a significant reduction in VC may be noted as indicated by other authors.<sup>2</sup> Various group pairing showed that there is significant difference in the TV, ERV and IC among the various groups of which some are statistically significant and comparable to other study.<sup>1</sup>

In a similar study, comparing lung function between singleton and twin pregnancy, no statistically significant differences were reported, which is comparable to our findings.<sup>7</sup> It is evident that the respiratory changes in pregnancy are largely mediated by hormonal factors, specially progesterone and to a lesser extent by estrogen. The mechanical effects of the over distended gravid uterus cause relatively little changes in the

**Table 1** Descriptive statistics of numeric variables of different groups

Variable	Nonpregnant (n = 20)		Early pregnancy (< 20 weeks) (n = 20)		Singleton pregnancy (36 weeks) (n = 20)		Multifetal gestation and/or polyhydramnios (36 weeks) (n = 20)		p-value (Kruskal-Wallis test)
	Mean (min-max)	SD	SE	Mean (min-max)	SD	SE	Mean (min-max)	SD	
<i>Gestational Age (days)</i>	—	—	—	75.90 (65-84)	5.49	1.22	252 (252-252)	2.17	0.46
<i>Age (Years)</i>	25.45 (18-31)	4.73	0.97	24.60 (18-31)	3.91	0.87	24.95 (21-31)	3.66 (18-30)	0.82
<i>Height (cm)</i>	150.35 (144-160)	5.05	1.13	151.20 (135-162)	7.35	1.64	153.15 (146-161)	4.24 (145-160)	0.95
<i>Weight (kg)</i>	53.45 (42-60)	4.18	0.93	50.30 (40-61)	5.31	1.18	60.20 (50-70)	8.64 (45-79)	1.89
<i>Blood pressure</i>									
<i>Systolic</i>	118.30 (110-140)	9.11	2.03	113 (110-130)	5.71	1.27	118.50 (110-140)	9.11 (110-140)	2.03
<i>Diastolic</i>	76 (60-70)	7.53	1.68	74.5 (60-80)	6.04	1.35	76 (60-90)	7.60 (60-80)	1.70
<i>Hb(gm/dl)</i>	11.53 (10.80-12.40)	0.44	0.09	11.48 (10.8-12.2)	0.56	0.08	11.50 (11-12.5)	0.40	0.09

**Table 2** Descriptive statistics of lung function tests of different groups

Variable	Nonpregnant (n = 20)		Early pregnancy (< 20 weeks) (n = 20)		Singleton pregnancy (36 weeks) (n = 20)		Multifetal gestation and/or polyhydramnios (36 weeks) (n = 20)		p-value (Kruskal-Wallis test)
	Mean (min-max)	SD	SE	Mean (min-max)	SD	SE	Mean (min-max)	SD	
<i>TV (Lt)</i>	0.56 (0.39-1.08)	0.18	0.04	0.73 (0.34-1.30)	0.24	0.05	0.88 (0.40-1.20)	0.21	0.002
<i>IRV (Lt)</i>	1.04 (0.53-1.80)	0.28	0.06	1.03 (0.54-1.78)	0.30	0.07	1.09 (0.26-1.65)	0.33	0.593
<i>ERV (Lt)</i>	0.76 (0.42-1.06)	0.15	0.03	0.74 (0.23-1.20)	0.31	0.07	0.49 (0.07-1.30)	0.29	0.000
<i>IC (Lt)</i>	1.61 (1.84-3.29)	0.41	0.09	1.65 (1.88-2.49)	0.39	0.09	1.95 (1.26-2.63)	0.35	0.016
<i>VC (Lt)</i>	2.35 (1.84-3.29)	0.36	0.08	2.44 (1.77-3.12)	0.41	0.09	2.44 (1.83-3.93)	0.46	0.921

TV = Tidal volume; IRV = Inspiratory reserve volume; ERV = Expiratory reserve volume; IC = Inspiratory capacity; VC = Vital capacity.

Group	Significance ( <i>p</i> -value $\leq 0.05$ is significant)		
	TV	ERV	IC
0-1	0.029	0.957	0.588
0-2	0.015	0.000	0.023
0-3	0.000	0.000	0.007
1-2	0.310	0.011	0.070
1-3	0.044	0.011	0.034
2-3	0.543	0.343	1.000

Nonpregnant = group 0; singleton early pregnancy = group 1; singleton pregnancy (36 weeks) = group 2; multifetal/polyhydramnios (36 weeks) = group 3  
 IC = inspiratory capacity; TV = tidal volume; ERV = expiratory reserve volume

pulmonary mechanics as it is compensated by increased circumference of thoracic cage as indicated by other studies<sup>2</sup> without any major change of respiratory muscle force mechanism.<sup>8</sup>

Though multifetal gestation is associated with higher levels of progesterone, more oxygen consumption, more demands by fetuses and uterine over distention, the lung volume changes are similar to that seen in the singleton pregnancy. As over distention as such contributes very little to the alteration of lung volumes polyhydramnios does not seem to cause significant respiratory compromise.

## CONCLUSION

It is evident from the study that a healthy woman without any respiratory or cardiovascular pathology can well-tolerate the

mechanical compression upon the diaphragm by the enlarging uterus and also the increasing demands of the growing fetus in multifetal pregnancy/polyhydramnios. It will help in better understanding of maternal respiratory physiology, and also management of the occasional cases of respiratory or cardiac disease with compromised respiratory function, where airway management and prevention of maternal hypoxemia are most important.

## REFERENCES

1. Elkus R, Popovich J Jr. Respiratory physiology in pregnancy. Clin Chest Medicine Dec 1992;13(4):555-65.
2. Madapp T, Sharma S. Pulmonary disease and pregnancy. Medline Search, Updated Oct 2010, emedicine.medscape.com/article.
3. Maternal physiological changes in pregnancy from Wikipedia. [http://en.wikipedia.org/wiki/Maternal\\_Physiological\\_Changes\\_in\\_Pregnancy](http://en.wikipedia.org/wiki/Maternal_Physiological_Changes_in_Pregnancy).
4. Christopher F, Ciliberto, Gertic F Marx. Physiological Changes Associated with pregnancy. Physiology 1998;9:1-3.
5. Gilroy RJ, Mangura BT, Levietes MH. Rib cage and abdominal volume displacement during breathing in pregnancy. Am Rev Respir Dis 1988;137(3):668-72.
6. Margareta M, Robb WG, Caren BD, Michael PH. Pulmonary gas exchange improves in the prone position with abdominal distention. Am J Respir Crit Care Med June 1988;157(6):1785-90.
7. McAuliffe F, Kametas N, Costello J, Rafferty GF, Greenough A, Nicolaidis K. Respiratory function in singleton and twin pregnancy. BJOG Jul 2002;109(7):765-69.
8. Lemos A, Caminla MA, Melo Jr EF, Domelas deAndrade A. Respiratory muscle force assessment in final 3 months of pregnancy. Braz J Phys Ther 2005;9(2):151-56.