

CASE REPORT

Successful Conservative Management of Spontaneous Unilateral Adrenal Hemorrhage in Pregnancy

¹Vineet V Mishra, ²Preeti Goyal, ³Sumesh Choudhary, ⁴Rohina Aggarwal, ⁵Priyankur Roy

ABSTRACT

Spontaneous adrenal hemorrhage (SAH) in the absence of trauma or adrenal tumor is a rare condition. Its incidence has been reported from 0.14 to 1.1% and it usually involves the right adrenal gland. During pregnancy, unilateral SAH has been reported very rarely. We describe a case who presented to us with pregnancy-induced hypertension and pain in right flank region for 1 day and diagnosed as a case of spontaneous right adrenal hemorrhage and managed conservatively.

Keywords: Adrenal hemorrhage, Pregnancy, Pregnancy-induced hypertension.

How to cite this article: Mishra VV, Goyal P, Choudhary S, Aggarwal R, Roy P. Successful Conservative Management of Spontaneous Unilateral Adrenal Hemorrhage in Pregnancy. *J South Asian Feder Obst Gynae* 2017;9(3):287-288.

Source of support: Nil

Conflict of interest: None

Date of received: 11 May 2017

Date of acceptance: 6 July 2017

Date of publication: August 2017

INTRODUCTION

Spontaneous adrenal hemorrhage (SAH) is an acute hemorrhage of the adrenal gland that occurs in the absence of prior trauma or adrenal tumor. The main symptoms are flank pain, hemorrhagic shock, and fever in some cases. The incidence of SAH has been reported from 0.14 to 1.1% and it usually involves the right gland.¹ Unilateral adrenal hemorrhage is infrequently associated with otherwise uncomplicated pregnancy, neurofibromatosis, and long-term nonsteroidal anti-inflammatory drug use.² During pregnancy, idiopathic unilateral SAH is reported to be a rare event with unknown incidence.³ We describe a case

who came to us with pregnancy-induced hypertension (PIH) and pain in right flank region for 1 day and was diagnosed to have spontaneous right adrenal hemorrhage with successful conservative management.

CASE REPORT

A 30-year-old female G3P2L2 with 32 weeks period of gestation and PIH presented to our department with pain abdomen in right flank region for 1 day. Her previous obstetric history was uneventful.

She was hemodynamically stable. On physical examination, her blood pressure was around 160/110 mm Hg, pulse rate was 92 per minute, and body temperature was 37.2°C. She had pitting edema in the lower limbs. She was on antihypertensive medications for last 15 days. Her obstetric examination revealed a relaxed uterus, regular fetal heart rate, and tenderness in right flank region. Her hemogram and serum biochemistry parameters were within normal limits. Abdominal ultrasound (ultrasonography) showed a single live pregnancy of 32 weeks with absent liquor and deranged color Doppler with brain-sparing effect parameters along with approximately 6.5 × 5.7 cm heterogeneous lesion in right lumbar region compatible with hemorrhage in the right adrenal gland. Magnetic resonance imaging (MRI) confirmed the diagnosis of adrenal hemorrhage.

Her cesarean was done after steroid coverage for the sake of baby. Patient's intrapartum and postpartum periods were uneventful except for the persistent right flank pain for which she was managed conservatively. She got discharged after 2 weeks and was followed up in outpatient department every 15 days. There were no complaints of flank pain after 1 month. Her repeat computed tomography (CT) scan revealed decreased size of adrenal hemorrhage.

DISCUSSION

Adrenal hemorrhage is a relatively uncommon condition with a variable and nonspecific presentation. The main symptoms of the condition are hemorrhagic shock, flank pain, and fever. Signs of acute abdomen including guarding, rigidity, or rebound tenderness have been reported in 15 to 20% because of the retroperitoneal location of adrenals.⁴

¹Professor and Head, ²Senior Resident, ³Assistant Professor
⁴Professor, ⁵Clinical Fellow

¹⁻⁵Department of Obstetrics and Gynecology, Smt. G.R. Doshi and Smt. K.M. Mehta Institute of Kidney Disease and Research Centre, Ahmedabad, Gujarat, India

Corresponding Author: Vineet V Mishra, Professor and Head Department of Obstetrics and Gynecology, Smt. G.R. Doshi and Smt. K.M. Mehta Institute of Kidney Disease and Research Centre, Ahmedabad, Gujarat, India, Phone: +917922687038 e-mail: vineet.mishra.ikdrc@gmail.com

Although the pathogenesis of adrenal hemorrhage is unclear, in nontraumatic cases, available evidence has implicated adrenocorticotrophic hormone, adrenal vein spasm, thrombosis, and limited venous drainage of adrenal as the mechanism behind the condition.⁴ Obstetric causes of bilateral adrenal hemorrhage include toxemia of pregnancy, spontaneous abortion, postpartum hemorrhage, twisted ovarian cyst in pregnancy, and more recently described antiphospholipid antibody syndrome.⁵ The present case had also PIH. Isolated case reports in association with long-term nonsteroidal anti-inflammatory drug use in uncomplicated pregnancy and neurofibromatosis have been reported.

Ultrasonography examination of the adrenals including Doppler study provides useful information regarding hemorrhage into the adrenal gland. Several weeks after the acute event, as the hematoma becomes cystic, the central echogenicity associated with adrenal hemorrhage decreases.

Contrast CT scan of the adrenals is the study of choice in demonstrating the adrenal hemorrhage. In pregnancy, MRI is the investigation of choice. The findings suggestive of hemorrhage into the adrenal are adrenal echogenicity with contrast enhancement, streaky appearance of perirenal fat, and perinephric hematoma. Several weeks after the acute hemorrhage, the CT scan shows a gradual decrease in size and attenuation.

Spontaneous unilateral adrenal hemorrhage during pregnancy has rarely been described. Our patient presented with right flank pain and PIH. Magnetic resonance imaging confirmed the diagnosis of unilateral adrenal hemorrhage. Patient was managed conservatively. Cesarean was done for fetal indication. On follow-up

after 1 month, patient's pain got resolved and CT showed resolving hemorrhage.

Spontaneous adrenal hemorrhage may occur in pregnancy in the absence of trauma or sepsis and can be managed conservatively if patient is hemodynamically stable. Thus, adrenal hemorrhage should be considered in the differential diagnosis of abdominal or flank pain with retroperitoneal hematoma in pregnancy.

CONCLUSION

Spontaneous adrenal hemorrhage during pregnancy is a rare condition. It should be considered in the differential diagnosis of pain abdomen in flank region in pregnant women. Though it can lead to acute adrenal crisis and hemorrhagic shock, it can be managed conservatively if patients' vitals are stable.

REFERENCES

1. Gavrilova-Jordan LP, Edmister WB, Farrell MA, Watson WJ. Spontaneous adrenal hemorrhage during pregnancy: a review of the literature and a case report of successful conservative management. *Obstet Gynecol Surv* 2005 Mar;60(3):191-195.
2. Merkle W. Spontaneous adrenal gland hemorrhage in adults. *Urologe A* 1986 Nov;25(6):343-346.
3. Kazarians VB, Kausch I, Gellifßen J, Doehn C, Jocham D. Spontaneous hemorrhage of the adrenal gland during pregnancy. *Aktuel Urol* 2007 Sep;38(5):403-405.
4. Kovacs KA, Lam YM, Pater JL. Bilateral massive adrenal hemorrhage. Assessment of putative risk factors by case controlled method. *Medicine (Baltimore)* 2001 Jan;80(1):45-53.
5. Espinosa G, Santos E, Cervera R, Piette JC, Asherson RA. Adrenal involvement in antiphospholipid syndrome: clinical and immunological characteristic of 86 patients. *Medicine (Baltimore)* 2003 Mar;82(2):106-118.